REFERENCES

1. Abhijit A. Date et. al., (2007), Design and evaluation of self-nanoemulsifying drug delivery systems (SNEDDS) for cefpodoxime proxetil, International Journal of Pharmaceutics, 329, 166–172.
2. Ahmed S. Zidan et. al., (2007), Quality by design: Understanding the formulation variables of a cyclosporine A self-nanoemulsified drug delivery systems by Box–Behnken design and desirability function, International Journal of Pharmaceutics, 332, 55–63.
3. Ana Maria Sierra Villar et. al., (2012), Design and optimization of self-nanoemulsifying drug delivery systems (SNEDDS) for enhanced dissolution of gemfibrozil, International Journal of Pharmaceutics, 431, 161– 175.
4. Anne T. Larsen et. al., (2013), Oral bioavailability of cinnarizine in dogs: Relation to SNEDDS droplet size, drug solubility and in vitro precipitation, European Journal of Pharmaceutical Sciences, 48, 339–350.
5. Arik Dahan et. al., (2008), Rationalizing the selection of oral lipid based drug delivery systems by an in vitro dynamic lipolysis model for improved oral bioavailability of poorly water soluble drugs, Journal of Controlled Release, 129, 1–10.
6. Basavaraj S et. al., (2014), Can formulation and drug delivery reduce attrition during
   drug discovery and development—review of feasibility, benefits and challenges, Acta Pharmaceutica Sinica B, 4(1), 3–17.
7. Dimitrios G. Fatouros, et. al., (2007), Morphological observations on a lipid-based drug delivery system during in vitro digestion, European Journal of Pharmaceutical Sciences, 31, 85–94.
8. Du Hyung Choi et. al., (2014), Evaluation of taste-masking effects of pharmaceutical sweeteners with an electronic tongue system, Drug Devlopment and Industrial Pharmacy, 40(3), 308–317.
9. Emad B. Basalious et. al.,, (2010), SNEDDS containing bioenhancers for improvement of dissolution and oral absorption of lacidipine. I: Development and optimization, International Journal of Pharmaceutics, 391, 203–211.

10. Flemming Seier Nielsen et. al., (2008), Bioavailability of probucol from lipid and surfactant based formulations in minipigs: Influence of droplet size and dietary state, European Journal of Pharmaceutics and Biopharmaceutics, 69, 553–562.
11. Gour Mukherji et. al., (2003), Taste masked composition, United states Patent, Patent No. US 6,565,877 B1, Patent date May 20, 2003.
12. Harini Chowdary Vadlamud et. al., (2016), Potential of microemulsified entacapone drug delivery systems in the management of acute Parkinson's disease, Journal of Acute Disease, 5(4), 315–325.
13. Huiyi Wu et. al., (2014), Combined use of phospholipid complexes and self-emulsifying microemulsions for improving the oral absorption of a BCS class IVcompound, baicalin, Acta Pharmaceutica Sinica B, 4(3), 217–226.
14. Hyeongmin Kin et. al., (2013), Liposomal formulations for enhanced lymphatic drug delivery, Asian Journal of Pharmaceutical Sciences, 8, 96-103.
15. Igor Legen et. al., (2013), Self-microemlsifying drug delivery system, United states Patent, Patent No. US8,592,490 B2, Patent date Nov. 26,2013.
16. JenniferWalsh et. al., Playing hide and seek with poorly tasting paediatric medicines: Do not forget the excipients, Advanced Drug Delivery Reviews, 73, 14–33.
17. Jing Lin, (2012), Self-microemulsifying dosage forms of low solubility active ingredients such as Co-enzyme Q10, United States Patent, Patent No. US 8,252,326 B2 , Patent date Aug. 28, 2012.
18. John W. Mauger et. al., (1998), Coating technology for taste masking orally administered bitter drugs, United states Patent, Patent No. 5,728,403, Patent date Mar. 17, 1998.
19. Kanchan Kohli et. al., (2014), Self emulsifying drug delivery system for a curcuminoid based composition, United States Patent, Patent No. US 8,835,509 B2, Patent date Sep. 16, 2014.
20. Katharina Woertz et. al., (2011), Development of a Taste-Masked Generic Ibuprofen Suspension: Top-Down Approach Guided by Electronic Tongue Measurements, Journal of Pharmaceutical Sciences, Vol. 100, No. 10, 4460 – 4470.

21. Kazunori Iwanaga et. al., (2015), Enhancing mechanism of intestinal absorption ofhighly lipophilic compounds using microemulsion - Quantitative analysis of the partitioning to the mesenteric lymph in intestinal cells, Asian Journal of Parmaceutical Scienses, 10, 186-193.
22. Lianli Li et. al., (2007), Utilization of a Modified Special-Cubic Design and an Electronic Tongue for Bitterness Masking Formulation Optimization, Journal of Pharmaceutical Sciences, Vol. 96, No. 10, 2723-2734.
23. Ling Wang et. al., (2015), Preparation and evaluation of taste masked oral suspension of arbidol hydrochloride, Asian Journa l of Pharmaceut ical Sciences, 10, 73 – 79.
24. Naser M.Y. Hasan, (2014), Role of medium-chain fatty acids in the emulsification mechanistics of self-micro-emulsifying lipid formulations, Saudi Pharmaceutical Journal, 22, 580–590.
25. Nicky Thomas et. al., (2013), Supersaturated Self-Nanoemulsifying Drug Delivery Systems (Super-SNEDDS) Enhance the Bioavailability of the Poorly Water-Soluble Drug Simvastatin in Dogs, The AAPS Journal, Vol. 15, No. 1, 219-227.
26. Orlagh M. Feeney et. al., (2016), 50 years of oral lipid-based formulations: Provenance, progress and future perspectives, Advanced Drug Delivery Reviews, 101, 167–194.
27. R. B. Shah et. al., (2007), Quality by design: Characterization of self-nano-emulsified drug delivery systems (SNEDDs) using ultrasonic resonator technology, International Journal of Pharmaceutics, 341, 189–194.
28. Sandeep Kalepu et. al., (2015), Insoluble drug delivery strategies: review of recent advances and business prospects, Acta Pharmaceutica Sinica B, 5(5), 442–453.
29. Hyeongmin Kim et. al., Liposomal formulations for enhanced lymphatic drug delivery, Asian Joural of Pharmaceutical Sciences, 8, 96 -103.
30. Sandeep Kalepu et. al., (2013), Oral lipid-based drug delivery systems – an overview, Acta Pharmaceutica Sinica B, 3(6), 361–372.
31. Shweta Gupta et. al., (2013), Formulation Strategies to Improve the Bioavailability of Poorly Absorbed Drugs with Special Emphasis on Self-Emulsifying Systems, ISRN Pharmaceutics, Volume 2013, 1 – 16.

32. Sivacharan Kollipara et. al., (2014), Pharmacokinetic aspects and in vitro–in vivocorrelation potential for lipid-based formulations, Acta Pharmaceutica Sinica B, 4(5), 333–349.
33. Suhua Li et. al., (2017), Effect of ionization of drug on drug solubilization in SMEDDS prepared using Capmul MCM and caprylic acid, Asian Journal of Pharmaceutical Sciences, 12, 73–82.
34. Sunil Kamboj et. al., (2016), Quality-by-design based development of a self-microemulsifying drug delivery system to reduce the effect of food on Nelfinavir mesylate, International Journal of Pharmaceutics, 501, 311–325.
35. Tamami Haraguchi et. al., (2016), Usefulness and limitations of taste sensors inthe evaluation of palatability and taste-masking in oral dosage forms, Asian Journal of Pharmaceutical Sciences, 11, 479–485.
36. Vijay D. Wagh et. al., (2009), Taste Masking Methods and Techniques in OralPharmaceuticals: Current Perspectives, Journal of Pharmacy Research, 2(6),1049-1054.
37. Yi Zhao et. al., (2010), Self-nanoemulsifying drug delivery system (SNEDDS) for oral delivery of Zedoary essential oil: Formulation and bioavailability studies, International Journal of Pharmaceutics, 383, 170–177.
38. Yoo-Jeong Jang et. al., (2016), Preparation and characterization of solidifiedSMEDDS containing flurbiprofen by spray drying method, Asian Journal of Pharmaceutical Sciences, 11, 108–109.
39. Yosra S.R. Elnaggar et. al., (2009), Self-nanoemulsifying drug delivery systems of tamoxifen citrate: Design and Optimization, International Journal of Pharmaceutics, 380, 133–141.
40. Yuqian Du et. al., (2013), Development and evaluation of taste-masked dry suspension of cefuroxime axetil for enhancement of oral bioavailability, Asian Journal of pharmaceutical sciences, 8, 287-294.